

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08209 - Goat Anti-MERTK Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MERTK, c-mer proto-oncogene tyrosine kinase, MER, MGC133349,

c-mer, MER receptor tyrosine kinase, STK kinase

Official Symbol: MERTK

Accession Number(s): NP_006334.2

Human GeneID(s): 10461

Immunogen

Peptide with sequence C-EVHDSKPHEGRY, from the internal region of the protein sequence according to NP_006334.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx. 125kDa band observed in lysates of cell lines THP-1 and A549 (calculated MW of 110.0kDa according to NP_006334.2). A 110kDa band was alternatively observed in some lysates. Recommended concentration: 2.5-3.5μg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB08209 optimised QC. Primary incubation 1 hour at room temperature.

Image A: THP-1 cell lysate at primary Ab concentration 3ug/ml. Image B: A549 cell lysate at primary Ab concentration 3.5ug/ml. (Loaded 35µg protein in RIPA buffer, per lane).

Detected by chemiluminescence.