

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB05934 - Goat Anti-MERTK Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MERTK, c-mer proto-oncogene tyrosine kinase, mer, C-MER, c-mer,

MER, MGC133349, RP38, MER receptor tyrosine kinase, STK kinase

Official Symbol: MERTK

Accession Number(s): NP_006334.2

Human GenelD(s): 10461

Immunogen

Peptide with sequence C-DSSEGSEVLM, from the C Terminus of the protein sequence according to NP_006334.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: Approx. 30kDa band observed in Human Brain, Human Ovary, Human Kidney, MOLT-4 and U937 lysates at 1ug of primary. Please note that the band we observe is different from the predicted size of approx. 115kDa according to NP_006334.


Currently we cannot find an explanation for this in the literature.

IHC: Paraffin embedded Human Adrenal Gland. Recommended concentration: 3.75µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB05934 (3.75µg/ml) staining of paraffin embedded Human Adrenal Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.