



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05879-T - Goat Anti-MURF1 / TRIM63 (N Term) Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: TRIM63, MURF1, MURF-1, RNF28, ring finger protein 28, IRF, SMRZ, MURF2, FLJ32380, iris ring finger protein, muscle specific ring finger protein 2, striated muscle RING zinc finger protein, tripartite motif-containing 63, muscle specific ring finger protein 1

Official Symbol: TRIM63

Accession Number(s): NP_115977.2

Human GeneID(s): [84676](#)

Immunogen

Peptide with sequence DYKSSLIQDGNPM-C, from the N Terminus of the protein sequence according to NP_115977.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


IHC: In paraffin embedded Human Skeletal Muscle shows strong staining in rare nuclei, and weak staining of muscle fibres in longitudinal section. Recommended concentration: 1.25µg/ml.

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of A549 cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Cow



EB05879 (1.25µg/ml) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB05879 Immunofluorescence analysis of paraformaldehyde fixed A549 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).