



Everest Biotech Ltd  
Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD, United Kingdom

[everestbiotech.com](http://everestbiotech.com)

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel +44 1869 238326

Fax +44 1869 238327

**Research Use Only. Not for diagnostic or therapeutic use.**

Storage: For long-term storage keep aliquots at -20°C. (Store no longer than 12 months at 4°C). Minimize freezing and thawing.

## EB06667 - Goat Anti-MYD88 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** myeloid differentiation primary response gene 88, MYD88D, myeloid differentiation primary response gene (88), MYD88

**Official Symbol:** MYD88

**Accession Number(s):** NP\_002459.1

**Human GeneID(s):** [4615](#)

**Non-Human GeneID(s):** 17874 (mouse), 301059 (rat)

### Immunogen

Peptide with sequence C-IKYKAMKKEFP, from the internal region of the protein sequence according to NP\_002459.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 33kDa band observed in HumanThymus lysates (calculated MW of 33.2kDa according to NP\_002459.1). Recommended concentration: 0.03-0.1µg/ml.


**IHC:** In paraffin embedded Human Tonsil shows staining of clusters of cells some of which are likely plasma cells. Recommended concentration, 4-6µg/ml.

**Enzyme immunoassay:** Sandwich-type ELISA with increasing amount of recombinant MYD88 captured by a rabbit antibody. Recommended reporter concentration: 5-10ug/ml


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human, Mouse, Rat, Dog



EB06667 staining (0.03 $\mu$ g/ml) of human thymus lysate (RIPA buffer, 35 $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



EB06667 (4 $\mu$ g/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining. These results could not be obtained after antigen retrieval at pH6 with this batch of antibody.



EB06667 (5 $\mu$ g/ml) as the reporter with EB002006 as the capture rabbit antibody (5 $\mu$ g/ml).