

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07938 - Goat Anti-Neurturin Antibody

Size: 100µg specific antibody in 200µl

Target Protein

Principal Names: NRTN, neurturin, NTN Official Symbol: NRTN Accession Number(s): NP_004549.1 Human GeneID(s): <u>4902</u>

Immunogen

Peptide with sequence C-DAHSRYHTVHELSARE, from the C Terminus of the protein sequence according to NP_004549.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 25kDa band observed in Human Heart lysates (calculated MW of 22.4kDa according to NP_004549.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour. **IHC:** Paraffin embedded Human Colon (Submucosal Plexus). Recommended

IHC: Paraffin embedded Human Colon (Submucosal Plexus). Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human Expected from sequence similarity: Human



| | 250kDa 150kDa 100kDa 75kDa |
|---|-------------------------------------|
| | 50kDa |
| | 37kDa |
| - | 25kDa 20kDa |
| | 15kDa |

EB07938 (0.1µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB07938 (5µg/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.