



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08500 - Goat Anti-NMNAT3 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** NMNAT3, nicotinamide nucleotide adenylyltransferase 3, PNAT-3, PNAT3, pyridine nucleotide adenylyltransferase 3

**Official Symbol:** NMNAT3

**Accession Number(s):** NP\_835471.1; NP\_001186976.1; NP\_001307440.1; NP\_001307442.1

**Human GeneID(s):** [349565](#)

### Immunogen

Peptide with sequence C-GSTWKGKSTQSTE, from the C Terminus of the protein sequence according to NP\_835471.1; NP\_001186976.1; NP\_001307440.1; NP\_001307442.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Preliminary testing showed a band at approx 30kDa in Human Placenta and Spleen lysate, and approx. 28kDa in Human Bone Marrow lysate after 0.3-0.5µg/ml antibody staining (calculated MW of 28.3kDa according to NP\_001307440.1). Primary incubation 1 hour at room temperature.

**Flow Cytometry:** A customer reported weak staining of HEK293 cells at a concentration of 10ug/ml.

### Species Reactivity

**Tested:**

**Expected from sequence similarity:** Human