



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05905 - Goat Anti-Npap60 / Nup50 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** NUP50, NPAP60, NPAP60L, MGC39961, DKFZP564A043, nucleoporin 50kDa, nuclear pore-associated protein 60L, CTA-268H5.7, OTTHUMP00000199530

**Official Symbol:** NUP50

**Accession Number(s):** NP\_009103.2; NP\_705931.1

**Human GeneID(s):** [10762](#)

**Non-Human GeneID(s):** 18141 (mouse)

**Important Comments:** This antibody is expected to recognise both reported isoforms (NP\_009103.2; NP\_705931.1).

### Immunogen

Peptide with sequence C-ELHKILLEKKDA, from the C Terminus of the protein sequence according to NP\_009103.2; NP\_705931.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.


**Western blot:** Approx 60kDa band observed in lysates of cell line Jurkat (calculated MW 50.1 kDa according to NP\_009103.2). Recommended for use at 0.2-1µg/ml.

**IHC:** In paraffin embedded Human Uterus shows textured nuclear staining of endometrial cells. Recommended concentration, 2.5-3.8µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Cow



EB05905 staining (0.5 $\mu$ g/ml) of Jurkat lysate (RIPA buffer, 30 $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.



EB05905 (3.8 $\mu$ g/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.