



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10571 - Goat Anti-NPAS4 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: NPAS4, neuronal PAS domain protein 4, Le-PAS, NXF, PASD10, bHLHe79, HLH-PAS transcription factor NXF, PAS domain-containing protein 10, class E basic helix-loop-helix protein 79, neuronal PAS4

Official Symbol: NPAS4

Accession Number(s): NP_849195.2

Human GeneID(s): [266743](#)

Non-Human GeneID(s): 225872 (mouse), 266734 (rat)

Immunogen

Peptide with sequence CRFNTSKSLRRQS, from the internal region of the protein sequence according to NP_849195.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.


Western blot: Approx 90kDa band observed in Human Brain (Frontal Cortex) lysates and in Mouse and Rat Brain lysates (calculated MW of 87.1kDa according to NP_849195.2).

Recommended concentration: 0.3-1µg/ml.


Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB10571 (0.3 μ g/ml) staining of Human Frontal Cortex lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB10571 (1 μ g/ml) staining of Mouse (A) and Rat (B) Brain lysates (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.