



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB12373 - Goat Anti-OAS2 (aa357-371) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** OAS2, 2'-5'-oligoadenylate synthetase 2, 69/71kDa, (2'-5')oligo(A) synthetase 2, 2'-5'-oligoadenylate synthase 2, 2-5A synthase 2, p69 OAS / p71 OAS

**Official Symbol:** OAS2

**Accession Number(s):** NP\_058197.2; NP\_002526.2

**Human GeneID(s):** [4939](#)

**Important Comments:** This antibody is expected to recognize reported isoform 1 (NP\_058197.2) and isoform 2 (NP\_002526.2).

### Immunogen

Peptide with sequence QPNKCFLEQIDSAVN, from the internal region of the protein sequence according to NP\_058197.2; NP\_002526.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.


**Western blot:** Approx 70kDa band observed in lysates of cell lines Daudi, Jurkat and K562 (calculated MW of 78.8kDa according to NP\_002526.2). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Lin et al, J Immunol. 2009 Dec 15;183(12):8035-43. PMID: 19923450). Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Breast. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB12373 (0.3 $\mu$ g/ml) staining of Daudi (A), Jurkat (B) and K562 (C) lysates (35 $\mu$ g protein in RIPA buffer).  
Detected by chemiluminescence.



EB12373 (5 $\mu$ g/ml) staining of paraffin embedded Human Breast. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.