

### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10833 - Goat Anti-Orexin Receptor 2 (aa405-419) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: HCRTR2, hypocretin (orexin) receptor 2, OX2R, hypocretin receptor

type 2, orexin receptor type 2, ox-2-R, ox2-R

Official Symbol: HCRTR2

Accession Number(s): NP\_001517.2

Human GenelD(s): 3062

Non-Human GenelD(s): 387285 (mouse), 25605 (rat)

## **Immunogen**

Peptide with sequence C-KSLTTQISNFDNISK, from the internal region (near C Terminus) of the protein sequence according to NP\_001517.2.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 50kDa band observed in Mouse fetal Brain lysates (calculated MW of 52.5kDa according to Mouse NP\_945200.1). Recommended concentration: 0.5-2µg/ml.

# **Species Reactivity**

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB10833 (0.5μg/ml) staining of Mouse fetal Brain lysate (35μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.