



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06152 - Goat Anti-ORP2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: OSBPL2, ORP2, ORP-2, MGC4307, MGC8342, FLJ20223, KIAA0772, oxysterol binding protein-like 2, OSBP-related protein 2, oxysterol-binding protein-related protein 2, oxysterol-binding protein-like protein 2

Official Symbol: OSBPL2

Accession Number(s): NP_055650.1; NP_653081.1

Human GeneID(s): [9885](#)

Non-Human GeneID(s): 228983 (mouse)

Important Comments: This antibody is expected to recognise both reported isoforms (represented by NP_055650.1 and NP_653081.1)

Immunogen

Peptide with sequence YFERNFSDCPDIY, from the C Terminus of the protein sequence according to NP_055650.1; NP_653081.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 55-60kDa band observed in Human Heart lysates (predicted MWs of 54 and 55kDa according to NP_055650.1 and NP_653081.1). Recommended for use at 1-2µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat



EB06152 staining (1.5µg/ml) of Human Heart lysate (RIPA buffer, 30µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.