



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06153 - Goat Anti-ORP3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: OSBPL3, ORP3, ORP-3, OSBP3, KIAA0704, MGC21526, oxysterol binding protein-like 3, OSBP-related protein 3, oxysterol-binding protein 3, oxysterol-binding protein-related protein 3, DKFZp667P1518, OTTHUMP00000122490, OTTHUMP00000122491, OTTHUMP00000202019, oxysterol-binding protein-like protein 3

Official Symbol: OSBPL3

Accession Number(s): NP_056365.1; NP_663160.1; NP_663161.1; NP_663162.1

Human GeneID(s): [26031](#)

Non-Human GeneID(s): 71720 (mouse)

Important Comments: This antibody is expected to recognise all reported isoforms (NP_056365.1; NP_663160.1; NP_663161.1; NP_663162.1).

Immunogen

Peptide with sequence MSDEKNLGVSQKL-C, from the N Terminus of the protein sequence according to NP_056365.1; NP_663160.1; NP_663161.1; NP_663162.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Immunofluorescence: Strong expression of the protein seen in the cytoplasm of A431 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB06153 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic and some nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06153 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.