



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.com

sales@everestbiotech.com

support@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

**Research Use Only. Not for
diagnostic or therapeutic use.**

Storage: For long-term storage
keep aliquots at -20°C. (Store no
longer than 12 months at 4°C).
Minimize freezing and thawing.

EB06228 - Goat Anti-PAR6alpha / PARD6A Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: partitioning-defective protein 6, partitioning defective-6 homolog alpha, par-6 partitioning defective 6 homolog alpha, Tax-interacting protein 40, PAR6C, PAR6, par-6 (partitioning defective 6, C.elegans) homolog alpha, Tax interaction protein 40, par-6 partitioning defective 6 homolog alpha (C.elegans), TIP-40, PAR-6A, TAX40, PAR-6, PAR6alpha, PARD6A

Official Symbol: PARD6A

Accession Number(s): NP_058644.1; NP_001032358.1

Human GeneID(s): [50855](#)

Important Comments: This antibody is expected to recognise both reported isoforms (NP_058644.1; NP_001032358.1).

Immunogen

Peptide with sequence C-GSRIRGDGSGFSL, from the C Terminus of the protein sequence according to NP_058644.1; NP_001032358.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.


Western blot: Preliminary experiments gave no signal but low background in Human Brain and Pancreas extracts at up to 1µg/ml.

IHC: In paraffin embedded Human Pancreas, shows strong cytoplasmic staining in rare cells of intralobular ducts. Recommended concentration, 10µg/ml. Also tested in paraffin embedded Human Heart where it shows speckled staining on myocardial fibres in transverse section. Recommended concentration, 5-10µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB06228 (10µg/ml) staining of paraffin embedded Human Pancreas. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.



EB06228 (5µg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.