

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford

Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB10154 - Goat Anti-Pex1 (mouse) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: 5430414H02Rik, E330005K07Rik, MGC102188, peroxin1, peroxisomal

biogenesis factor 1, peroxisome biogenesis disorder protein 1, ZWS1, Pex1

Official Symbol: Pex1

Accession Number(s): NP\_082053.1

Non-Human GenelD(s): 71382 (mouse), 500006 (rat)

## **Immunogen**

Peptide with sequence C-QAKESTFPKEGDAH, from the internal region of the protein

sequence according to NP\_082053.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 150kDa band observed in Mouse and Rat Liver lysates (calculated MW of 137kDa according to NP\_082053.1). Recommended concentration: 0.3-1µg/ml.

### **Species Reactivity**

Tested: Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa

EB10154 (0.3 $\mu$ g/ml) staining of Mouse Liver lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemilluminescence.

20kDa