

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07176 - Goat Anti-PIAS2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PIAS2, protein inhibitor of activated STAT, 2, MGC102682, MIZ1, PIASX-ALPHA, PIASX-BETA, SIZ2, miz, Msx-interacting-zinc finger, protein inhibitor of activated STAT X, ARIP3 Official Symbol: PIAS2 Accession Number(s): NP_775298.1; NP_004662.2 Human GeneID(s): 9063 Non-Human GeneID(s): 17344 (mouse) Important Comments: This antibody is expected to recognise both reported isoforms (NP_775298.1 and NP_004662.2). No cross-reactivity expected with PIAS1, 3 and 4.

Immunogen

Peptide with sequence KEAMKVSSQPCTKIE, from the internal region of the protein sequence according to NP_775298.1; NP_004662.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Preliminary experiments gave an approx 90kDa band in lysates of cell line A431after 0.5µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 68.2kDa according to NP_004662.2. The 90kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

IHC: In paraffin embedded Human Testis shows nuclear and extra-nuclear staining in spermatids. Recommended concentration, 3-5µg/ml.

Species Reactivity

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07176 (3.8 μ g/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.