

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06239 - Goat Anti-PIST / FIG / GOPC Antibody

Size: 100µg specific antibody in 200µl

Plite

Target Protein

Principal Names: PIST, GOPC, CAL, PDZ/coiled-coil domain binding partner for the rho-family GTPase TC10, fused in glioblastoma, CFTR-associated ligand, Golgi associated PDZ and coiled-coil motif containing protein, golgi associated PDZ and coiled-coil motif containing, CAL, FIG, GOPC1, dJ94G16.2, dJ94G16.2 PIST **Official Symbol:** GOPC

Accession Number(s): NP_065132.1; NP_001017408.1 Human GenelD(s): <u>57120</u>

Important Comments: This antibody is expected to recognize both reported isoforms (NP_065132.1; NP_001017408.1).

Immunogen

Peptide with sequence C-LDDLHTLYHKKSY, from the C Terminus of the protein sequence according to NP_065132.1; NP_001017408.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx. 55kDa band observed in Human Brain (Frontal Cortex) lysates (calculated MW of 50.5kDa according to NP_065132.1). Recommended concentration: 0.05-0.2µg/ml.

Immunofluorescence: This product has been successfully used on HeLa cells.

Species Reactivity

Tested: Human Expected from sequence similarity: Human, Dog, Cow

250kDa 150kDa 100kDa 75kDa 50kDa
37kDa
25kDa
20kDa
37kDa 25kDa

15kDa

EB06239 (0.05µg/ml) staining of Human Frontal Cortex lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.