



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB11960 - Goat Anti-PLA2G7 (aa53-64) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PLA2G7, phospholipase A2, group VII (platelet-activating factor acetylhydrolase, plasma), LDL-PLA2, LP-PLA2, PAFAD, PAFAH, 1-alkyl-2-acetyl-glycerophosphocholine esterase, 2-acetyl-1-alkyl-glycerophosphocholine esterase, LDL-PLA(2), LDL-associated phospholipase A2, PAF 2-acylhydrolase, PAF acetylhydrolase, gVIIA-PLA2, group-VIIA phospholipase A2, lipoprotein-associated phospholipase A2, platelet-activating factor acetylhydrolase

Official Symbol: PLA2G7

Accession Number(s): NP_005075.3

Human GeneID(s): [7941](#)

Important Comments: Reported variants represent identical protein: NP_005075.3, NP_001161829.1

Immunogen

Peptide with sequence C-QTKIPRGNGPYS, from the internal region of the protein sequence according to NP_005075.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Preliminary experiments gave an approx 100kDa band in Human Liver lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 50.1kDa according to NP_005075.3. The 100kDa band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Does the observed molecular weight correspond to a glycosylated form?

Species Reactivity

Tested:

Expected from sequence similarity: Human