



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10590 - Goat Anti-POLDIP2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: POLDIP2, polymerase (DNA-directed), delta interacting protein 2, DKFZp586F1524, PDIP38, POLD4, 38 kDa DNA polymerase delta interaction protein, p38, polymerase delta interacting protein 38, polymerase delta-interacting protein 2

Official Symbol: POLDIP2

Accession Number(s): NP_056399.1

Human GeneID(s): [26073](#)

Non-Human GeneID(s): 67811 (mouse)

Immunogen

Peptide with sequence C-RFERPDGSHFDVR, from the internal region (near C terminus) of the protein sequence according to NP_056399.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 37kDa band observed in Rat Vascular Smooth Muscle Cells lysates (calculated MW of 15.2kDa according to Rat NP_001099286.1). The band diminished upon siRNA knock down. Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Rat

Expected from sequence similarity: Human, Mouse, Dog, Cow



EB10590 (0.5µg/ml) staining of Rat VSMC lysate (35µg protein in RIPA buffer). Primary incubation was overnight at 4C. Detected by chemiluminescence. Data provided by Everest Grant winner Dr. B. Lassegue, Emory University, Atlanta, USA