



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09674 - Goat Anti-PPIG Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PPIG, peptidylprolyl isomerase G (cyclophilin G), CARS-Cyp, CYP, MGC133241, SRCyp, Clk-associating RS-cyclophilin, peptidyl-prolyl isomerase G (cyclophilin G), peptidylprolyl isomerase G

Official Symbol: PPIG

Accession Number(s): NP_004783.2

Human GeneID(s): [9360](#)

Non-Human GeneID(s): 228005 (mouse), 83624 (rat)

Immunogen

Peptide with sequence C-RDHNSSNNSREKKAD, from the internal region of the protein sequence according to NP_004783.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:1000.

Western blot: Approx 85kDa band observed in Human Kidney lysates (calculated MW of 88.6kDa according to NP_004783.2). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB09674 (2 μ g/ml) staining of Human Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.