

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07297 - Goat Anti-Prion Protein (143-153) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PRNP, prion protein (p27-30) (Creutzfeldt-Jakob disease, Gerstmann-Strausler-Scheinker syndrome, fatal familial insomnia), ASCR, CD230, CJD, GSS, MGC26679, PRIP, PrP, PrP27-30, PrP33-35C, PrPc, CD230 antigen, major prion protein, prion protein, prion protein (p27-30) (Creutzfeld-Jakob disease Gerstmann-Strausler-Scheinker syndrome, fatal familial insomnia), prion protein PrP,

prion-related protein

Official Symbol: PRNP

Accession Number(s): NP_000302.1

Human GenelD(s): 5621

Non-Human GenelD(s): 19122 (mouse), 24686 (rat)

Important Comments: Reported variants represent identical protein (NP_000302.1;

NP_898902.1; NP_001073590.1; NP_001073591.1; NP_001073592.1).

Immunogen

Peptide with sequence C-SDYEDRYYREN, from the internal region of the protein sequence according to NP_000302.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 25kDa band observed in Human Brain lysates (calculated MW of 27.7kDa according to NP_000302.1 and NP_898902.1). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Pig, Cow



EB07297 (0.3 μ g/ml) staining of Human Brain lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.