



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12886 - Goat Anti-PRKD1 (aa383-395) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: PRKD1, PKC-MU, PKCM, PKD, PRKCM;

Official Symbol: PRKD1

Accession Number(s): NP_002733.2

Human GeneID(s): [5587](#)

Immunogen

Peptide with sequence C-QDPDPDHEDANRT, from the internal region of the protein sequence according to NP_002733.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.


Western blot: In transfected HEK293 transiently expressing Human PKD1 (DYKDDDDK tagged), a band of approx. 100kDa is observed. The same band is observed using anti-DYKDDDDK antibody and no bands are observed in cells transiently expressing Human PKD2 or PKD3. Recommended concentration 0.1-0.3ug/ml

Immunoprecipitation: PKD1 was immunoprecipitated from Hek293T cells (as above). Immunoprecipitates were analyzed by Western Blot probed with the same antibody.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



HEK293 lysate overexpressing Human DYKDDDDK-tagged PKD1, Human DYKDDDDK-tagged PKD2 or Human GST-tagged PKD3 probed with EB12886 (0.1ug/ml) in top panel, probed with anti-DYKDDDDK in middle panel and probed with anti-GST in bottom panel. Data kindly obtained from Dr Peter Storz, Mayo Clinic, USA.



HEK293 lysate overexpressing Human DYKDDDDK-tagged PKD1 was used to immunoprecipitate PKD1 with 2ug EB12886. The precipitate was subsequently probed in Western blot using EB12886 at 1ug/ml. The secondary anti-goat picks up the heavy chain of EB12886 used for the immunoprecipitation (annotated as IgG). Data kindly obtained from Dr Peter Storz, Mayo Clinic, USA