



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09534 - Goat Anti-PRMT5 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PRMT5, protein arginine methyltransferase 5, HRMT1L5, IBP72, JBP1, SKB1, SKB1Hs, HMT1 hnRNP methyltransferase-like 5, SKB1 homolog

**Official Symbol:** PRMT5

**Accession Number(s):** NP\_006100.2; NP\_001034708.1

**Human GeneID(s):** [10419](#)

**Non-Human GeneID(s):** 27374 (mouse), 364382 (rat)

**Important Comments:** This antibody is expected to recognize both isoforms (NP\_006100.2; NP\_001034708.1).

### Immunogen

Peptide with sequence RACREKDRDPEAQ, from the internal region of the protein sequence according to NP\_006100.2; NP\_001034708.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 70kDa band observed in Jurkat lysates (calculated MW of 72.7kDa and 71.3kDa according to NP\_006100.2; NP\_001034708.1). Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Pig, Dog, Cow



EB09534 staining (2 $\mu$ g/ml) of Jurkat lysate (RIPA buffer, 35 $\mu$ g total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.