



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08195 - Goat Anti-Proenkephalin Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** proenkephalin, PENK

**Official Symbol:** PENK

**Accession Number(s):** NP\_006202.1

**Human GeneID(s):** [5179](#)

### Immunogen

Peptide with sequence C-RSHHQDGSNDNEE, from the internal region of the protein sequence according to NP\_006202.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 37kDa band observed in Human Adrenal Gland lysates (calculated MW of 30.8kDa according to NP\_006202.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Normant and Loh, Endocrinology. 1998 Apr;139(4):2137-45; PMID: 9529003). Recommended concentration: 1-3µg/ml.

**IHC:** Frozen section of Human Hypothalamus shows staining of dense enkephalinergic axon plexus and scattered neuronal cell bodies (higher magnification inset) in the human infundibular nucleus. Recommended concentration: 0.3-1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human

### Specific Reference

**This antibody has been successfully used in IHC in Human:**

Skrapits K, Borsay B, Herczeg L, Ciofi P, Liposits Z, Hrabovszky E.


Neuropeptide co-expression in hypothalamic kisspeptin neurons of laboratory animals and the human.

Front Neurosci. 2015 Feb 10;9:29.

PMID: 25713511



EB08195 (0.3 $\mu$ g/ml) staining of Human Adrenal Gland lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB08195 (0.3 $\mu$ g/ml) staining of PFA-perfused cryosection of Human Hypothalamus. Antigen retrieval with citrate buffer pH 6 at 80C for 30min, Cy3-staining. Data obtained by Prof. Erik Hrabovszky, Inst, Exp, Med., Budapest, Hungary.