



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05535 - Goat Anti-RANBP8 / IPO8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: IPO8, importin 8, RANBP8, RAN-binding protein 8, FLJ26580, RAN binding protein 8

Official Symbol: IPO8

Accession Number(s): NP_006381.2; NP_001177924.1

Human GeneID(s): [10526](#)

Important Comments: This antibody is expected to recognise isoform 1 (NP_006381.2) and isoform 2 (NP_001177924.1).

Immunogen

Peptide with sequence C-SAFNFGTVPSNN, from the C Terminus of the protein sequence according to NP_006381.2; NP_001177924.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Preliminary experiments in lysates of cell line HeLa gave no specific signal but low background (at antibody concentration up to 1µg/ml).

IHC: In paraffin embedded Human Tonsil shows cytoplasm / perinuclear staining in the parenchyma. Recommended concentration, 4-6µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog, Pig, Cow



EB05535 (4µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining. Similar results were obtained after antigen retrieval at pH9.