



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.com

sales@everestbiotech.com

support@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

Research Use Only. Not for diagnostic or therapeutic use.

Storage: For long-term storage keep aliquots at -20°C. (Store no longer than 12 months at 4°C). Minimize freezing and thawing.

EB05535 - Goat Anti-RANBP8 / IPO8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: RAN binding protein 8, FLJ26580, RAN-binding protein 8, RANBP8, importin 8, IPO8

Official Symbol: IPO8

Accession Number(s): NP_006381.2; NP_001177924.1

Human GeneID(s): [10526](#)

Important Comments: This antibody is expected to recognise isoform 1 (NP_006381.2) and isoform 2 (NP_001177924.1).

Immunogen

Peptide with sequence C-SAFNFGTVPSNN, from the C Terminus of the protein sequence according to NP_006381.2; NP_001177924.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Preliminary experiments in lysates of cell line HeLa gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates?

IHC: In paraffin embedded Human Tonsil shows cytoplasm staining in the parenchyma. Recommended concentration, 4-6µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog, Pig, Cow



EB05535 (4 μ g/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining. Similar results were obtained after antigen retrieval at pH9.