

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

Enquiries:

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06543 - Goat Anti-RIC8A / Synembryn Antibody

Size: 100µg specific antibody in 200µl

P_{lite}

Target Protein

Principal Names: RIC8A, synembryn, resistance to inhibitors of cholinesterase 8 homolog A (C. elegans), MGC104517, MGC131931, RIC8, RIC8 protein, resistance to

inhibitors of cholinesterase 8 homolog A

Official Symbol: RIC8A

Accession Number(s): NP_068751.4

Human GenelD(s): 60626

Non-Human GenelD(s): 101489 (mouse)

Immunogen

Peptide with sequence CLSSDPDSDPD, from the C Terminus of the protein sequence according to NP_068751.4.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 70kDa band observed in human brain lysates (calculated MW of 60.4kDa according to NP_068751.4). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Cow

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB06543 staining (1µg/ml) of human brain lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by western blot using chemiluminescence.