



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.com

sales@everestbiotech.com

support@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

**Research Use Only. Not for
diagnostic or therapeutic use.**

Storage: For long-term storage
keep aliquots at -20°C. (Store no
longer than 12 months at 4°C).
Minimize freezing and thawing.

EB05742 - Goat Anti-SART1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: small nuclear ribonucleoprotein 110kDa (U4/U6.U5), SNRNP110, squamous cell carcinoma antigen recognized by T cells 1, squamous cell carcinoma antigen recognised by T cells, U4/U6.U5 tri-snRNP-associated 110 kDa protein, SART1(800) protein, SART1(259) protein, IgE autoantigen, Snu66, SART1259, MGC2038, HOMS1, Ara1, ARA1, squamous cell carcinoma antigen recognised by T cells, SART1

Official Symbol: SART1

Accession Number(s): NP_005137.1

Human GeneID(s): [9092](#)

Non-Human GeneID(s): 20227 (mouse), 29678 (rat)

Important Comments: This antibody is expected to recognise both the human SART1(800) and SART1(259) proteins.

Immunogen

Peptide with sequence GSSKKHRGEKEAA-C, from the N Terminus of the protein sequence according to NP_005137.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 125kDa band observed in HeLa lysates (consistent with Vertegaal et al, below, calculated MW of 90.3 kDa according to NP_005137). Recommended concentration: 0.03-0.1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Cow



EB05742 (0.03 μ g/ml) staining of HeLa lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.