

## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11236 - Goat Anti-Slc5a1 (rat) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** high affinity sodium-glucose cotransporter, MGC93553, Na(+)/glucose cotransporter 1, SglT1, sodium/glucose cotransporter 1, solute carrier family 5 (sodium/glucose cotransporter), member 1, solute carrier family 5, member 1, Solute carrier family 5, member alpha 1 (Na+/glucose cotransporter), Slc5a1

**Official Symbol:** Slc5a1

**Accession Number(s):** NP\_037165.1

**Non-Human GeneID(s):** 25552 (rat)

### Immunogen

Peptide with sequence C-KDTIEIDAEAPQKEK, from the internal region of the protein sequence according to NP\_037165.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.


**Western blot:** Approx 70kDa band observed in Rat Kidney lysates (calculated MW of 72.1kDa according to NP\_037165.1). Recommended concentration: 0.3-2µg/ml. Primary incubation 1 hour at room temperature.

**IHC:** Paraffin embedded Rat Kidney. Recommended concentration: 10µg/ml.


### Species Reactivity

**Tested:** Rat


**Expected from sequence similarity:** Rat



EB11236 (1µg/ml) staining of Rat Kidney lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB11236 (10µg/ml): Staining of paraffin embedded Rat Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-Staining (1:500).



EB11236 Negative Control showing staining of paraffin embedded Rat Kidney, with no primary antibody.