

#### **UK Office**

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

# **US Office**

#### Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09792 - Goat Anti-SNX1 Antibody

Size: 100µg specific antibody in 200µl

# *Plus* lite

#### **Target Protein**

Principal Names: HsT17379, MGC8664, SNX1A, sorting nexin 1, sorting nexin 1A, Vps5, SNX1 Official Symbol: SNX1

Accession Number(s): NP\_003090.2; NP\_690039.1 Human GeneID(s): 6642 Non-Human GeneID(s): 56440 (mouse), 84471 (rat) Important Comments: This antibody is expected to recognize isoform a and c (NP\_003090.2; NP\_690039.1.

#### Immunogen

Peptide with sequence C-TNSSKPQPTYEELE, from the internal region of the protein sequence according to NP\_003090.2; NP\_690039.1.

Please note the <u>peptide</u> is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:64000.


**Western blot:** Approx 75kDa band observed in lysates of cell lines HeLa and U2OS (calculated MW of 59.1kDa according to NP\_003090.2). This molecular weight is routinely observed by other sources . Recommended concentration: 0.1-1µg/ml. Primary incubation 1 hour at room temperature.

**IHC:** In paraffin embedded Human Uterus shows vesicular staining in the cytoplasm of endometrial epithelium. Recommended concentration, 3-5µg/ml.


**Immunocytochemistry:** Vesicular staining of SNX1 (green) in formaldehyde-fixed SH5Y5Y. Recommended concentration, 3-5µg/ml.

#### **Species Reactivity**


Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB09792 (1µg/ml) staining of HeLa (A) and (0.1ug/ml) U2OS (B) cell lysate (35µg protein in RIPA buffer).. Detected by chemiluminescence.



EB09792 (3.8µg/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB09792 (5ug/ml) staining of formaldehyde-fixed SH5Y5Y and detected with Alexa 488 in confocal microscopy. Data obtained from Dr. Schallburg Nielsen, Aarhus University Denmark.