



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB09775 - Goat Anti-SRGAP2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** SRGAP2, FNBP2, formin binding protein 2, KIAA0456, SLIT-ROBO  
Rho GTPase activating protein 2, srGAP3

**Official Symbol:** SRGAP2

**Accession Number(s):** NP\_056141.2

**Human GeneID(s):** [23380](#)

**Non-Human GeneID(s):** 14270 (mouse), 360840 (rat)

**Important Comments:** This antibody is expected to recognize isoform a (NP\_056141.2).

### Immunogen

Peptide with sequence C-KQEDRQTPRSPDST, from the internal region of the protein sequence according to NP\_056141.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:8000.


**Western blot:** Approx 124kDa band observed in lysates of cell line HeLa and in rat primary culture cortical neurons at Day 4 (calculated MW of 121kDa according to human NP\_056141.2 but the rat protein sequence is still provisional). Recommended concentration: 0.1-0.3µg/ml.

**Immunofluorescence:** Transfected HEK293 transiently expressing GFP-tagged Human srGAP2 with the primary antibody stained in the red channel (Confocal IF).


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Cow



EB09775 (0.1 $\mu$ g/ml) staining of Rat Cortical Neuron lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence. Data kindly provided by Ms. Ya-Jing Mi and Dr. Wei-Lin Jin, Institute of Neurosciences, Shanghai Jiao Tong University.



HEK293 overexpressing Human srGAP2 and probed with EB09775 at 2.5 $\mu$ g/ml in the right panel. Data kindly provided by Ms. Ya-Jing Mi and Dr. Wei-Lin Jin, Institute of Neurosciences, Shanghai Jiao Tong University.