



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05885-B - Goat Anti-SSP29 / ANP32B, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ANP32B, acidic nuclear phosphoprotein 32 family member B, APRIL, PHAPI2, SSP29, acidic (leucine-rich) nuclear phosphoprotein 32 family, member B, acidic protein rich in leucines, putative HLA-DR-associated protein I-2, silver-stainable protein SSP29

Official Symbol: ANP32B

Accession Number(s): NP_006392.1

Human GeneID(s): [10541](#)

Non-Human GeneID(s): 67628 (mouse), 170724 (rat)

Important Comments: This antibody does NOT recognize the TNF family member also known as APRIL - Gene ID number 8741. However this antibody may cross-react with ANP32A (GeneID 8125).

Immunogen

Peptide with sequence C-KRKRETDDEGEDD., from the C Terminus of the protein sequence according to NP_006392.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 28kDa band observed in Human Tonsil lysates (calculated MW of 28.8kDa according to NP_006392.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat, Pig

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



Biotinylated EB05885 (0.2 μ g/ml) staining of Human Tonsil lysate (35 μ g protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.