

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07469 - Goat Anti-STIM1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: STIM1, stromal interaction molecule 1, D11S4896E, GOK

Official Symbol: STIM1

Accession Number(s): NP_003147.2; NP_001264890.1; NP_001264891.1

Human GeneID(s): [6786](#)

Non-Human GeneID(s): 20866 (mouse), 361618 (rat)

Immunogen

Peptide with sequence C-YIQNRYSKEHMKK, from the internal region of the protein sequence according to NP_003147.2; NP_001264890.1; NP_001264891.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Successful use of this antibody was reported in Western blot showing an approx. 80kDa in Human platelets: PMID 18439569

Flow Cytometry: Flow cytometric analysis of HepG2 cells. Recommended concentration: 10ug/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific Reference


This antibody has been successfully used in WB on Human Platelets (80kDa) in the following paper:

López JJ, Jardín I, Bobe R, Pariente JA, Enouf J, Salido GM, Rosado JA.

STIM1 regulates acidic Ca²⁺ store refilling by interaction with SERCA3 in human platelets.

Biochem Pharmacol. 2008 Jun 1;75(11):2157-64.

PMID: 18439569



EB07469 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.