

UK Office

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06180 - Goat Anti-Syntrophin (alpha 1) Antibody

Size: 100µg specific antibody in 200µl

C

Target Protein

Principal Names: SNTA1, syntrophin, alpha 1 (dystrophin-associated protein A1, 59kDa, acidic component), SNT1, TACIP1, dJ1187J4.5, pro-TGF-alpha cytoplasmic domain-interacting protein 1, dystrophin-associated protein A1, 59kD, acidic component, syntrophin, alpha 1 (dystrophin-associated protein A1, 59kD, acidic component), LQT12, acidic alpha 1 syntrophin Official Symbol: SNTA1 Accession Number(s): NP_003089.1 Human GeneID(s): <u>6640</u> Non-Human GeneID(s): 20648 (mouse)

Immunogen

Peptide with sequence ASGRRAPRTGLLE-C, from the N Terminus of the protein sequence according to NP_003089.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000. **Western blot:** Approx 60kDa band observed in Human and Mouse Muscle lysates and in rodent Heart lysates (calculated MW of 53.9kDa according to Human NP_003089.1). Recommended concentration: 1-3μg/ml.

Species Reactivity

Tested: Human, Mouse, Rat Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB06180 (0.01µg/ml) staining of Human Muscle lysate (35µg protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Primary incubation was 1 hour. Detected by chemiluminescence.