



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08872 - Goat Anti-TAP1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TAP1, transporter 1, ATP-binding cassette, sub-family B (MDR/TAP), ABC17, ABCB2, APT1, D6S114E, FLJ26666, FLJ41500, PSF1, RING4, TAP1*0102N, TAP1N, ABC transporter, MHC 1, ATP-binding cassette transporter, ATP-binding cassette, sub-family B (MDR/TAP), member 2, ATP-binding cassette, sub-family B, member 2, antigen peptide transporter 1, peptide supply factor 1, transporter 1, ATP-binding cassette, sub-family B, transporter associated with antigen processing, transporter, ATP-binding cassette, major histocompatibility complex, 1

Official Symbol: TAP1

Accession Number(s): NP_000584.2

Human GeneID(s): [6890](#)

Immunogen

Peptide with sequence C-QKFREKLQEIKT, from the internal region of the protein sequence according to NP_000584.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 85kDa band observed in lysates of cell line MCF7 (calculated MW of 87.2kDa according to NP_000584.2). Recommended concentration: 1-3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB08872 (1 μ g/ml) staining of MCF7 lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.