



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08727 - Goat Anti-TCF4 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TCF4, transcription factor 4, E2-2, ITF2, MGC149723, MGC149724, PTHS, SEF2, SEF2-1, SEF2-1A, SEF2-1B, transcription factor-4 (immunoglobulin transcription factor-2)

Official Symbol: TCF4

Accession Number(s): NP_001077431.1 ; NP_003190.1

Human GeneID(s): [6925](#)

Important Comments: This antibody is expected to recognise the reported isoforms a and b (NP_001077431.1; NP_003190.1)

Immunogen

Peptide with sequence C-NGGHPSRNYGDGT, from the internal region of the protein sequence according to NP_001077431.1 ; NP_003190.1 .

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 70kDa band observed in Human Heart lysates (calculated MW of 71.3kDa according to NP_003190.1). Recommended concentration: 0.5-1.5µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Dog



EB08727 (0.5µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.