



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB10507 - Goat Anti-TGM7 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TGMZ, transglutaminase 7, transglutaminase Z, TGM7

Official Symbol: TGM7

Accession Number(s): NP_443187.1

Human GeneID(s): [116179](#)

Non-Human GeneID(s): 640543 (mouse)

Immunogen

Peptide with sequence C-KIEISQQGHSVT, from the internal region of the protein sequence according to NP_443187.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 75kDa band observed in Human Lung and Testis and in Mouse Kidney, Lung and Testis lysates (calculated MW of 79.9kDa according to Human NP_443187.1 and 79.8kDa according to Mouse NP_001153896.1). Recommended concentration: 0.1-0.3µg/ml. An additional band of unknown identity was also consistently observed at 17kDa. This band was successfully blocked by incubation with the immunizing peptide.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Dog, Pig



EB10507 (0.1 μ g/ml) staining of Mouse Kidney lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB10507 (5 μ g/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.