

#### **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07967 - Goat Anti-THRB Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** THRB, thyroid hormone receptor, beta (erythroblastic leukemia viral (v-erb-a) oncogene homolog 2, avian), ERBA-BETA, ERBA2, GRTH, MGC126109, MGC126110, NR1A2, THR1, THRB1, THRB2, avian erythroblastic leukemia viral (v-erb-a) oncogene homolog 2, generalized resistance to thyroid hormone, oncogene ERBA2, thyroid hormone receptor beta 1, thyroid hormone receptor, beta

Official Symbol: THRB

Accession Number(s): NP\_000452.2

Human GenelD(s): 7068

Non-Human GenelD(s): 21834 (mouse), 24831 (rat)

**Important Comments:** This antibody is not expected to cross-react to THRA.

## **Immunogen**

Peptide with sequence C-REELQKSIGHKP, from the internal region of the protein sequence according to NP\_000452.2.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 50+60kDa bands observed in Human Bone Marrow lysates (calculated MW of 52.8kDa according to NP\_NP\_000452.2 and of 54.5kDa according to EAW64350.1). Recommended concentration: 0.3-1ug/ml

# **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07967 (0.3 $\mu$ g/ml) staining of Human Bone Marrow lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.