

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB12135 - Goat Anti-thromboxane A2 receptor (isoform beta) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TBXA2R, thromboxane A2 receptor, BDPLT13, TXA2-R, prostanoid TP

receptor

Official Symbol: TBXA2R

Accession Number(s): NP_963998.2

Human GenelD(s): 6915

Important Comments: This antibody is expected to recognize the reported isoform beta

(NP_963998.2) only.

Immunogen

Peptide with sequence C-PFEPPTGKALSRKD, from the C Terminus of the protein sequence according to NP_963998.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 55kDa band observed in Human Lung and Placenta lysates (calculated MW of 44.2kDa according to NP_963998.2). The observed molecular weight corresponds to the glycosylated form. Recommended concentration: 0.03-0.1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

EB12135 ($0.03\mu g/ml$) staining of Human Lung lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.