



Everest Biotech Ltd
Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD, United Kingdom

everestbiotech.com

sales@everestbiotech.com

support@everestbiotech.com

Tel +44 1869 238326

Fax +44 1869 238327

**Research Use Only. Not for
diagnostic or therapeutic use.**

Storage: For long-term storage
keep aliquots at -20°C. (Store no
longer than 12 months at 4°C).
Minimize freezing and thawing.

EB06323 - Goat Anti-TIP30 / CC3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: short chain dehydrogenase/reductase family 44U, member 1, SDR44U1, FLJ26963, HIV-1 Tat interactive protein 2, 30 kDa, HIV-1 Tat interactive protein 2, 30 kD, Tat-interacting protein (30kD), HIV-1 Tat interactive protein 2, 30kDa, HTATIP2, TIP30, CC3

Official Symbol: HTATIP2

Accession Number(s): NP_001091990.1; NP_006401.3; NP_001091991.1;
NP_001091992.1; NP_001091993.1

Human GeneID(s): [10553](#)

Important Comments: This antibody is expected to recognise all three reported isoforms. Variants (NP_006401.3; NP_001091991.1; NP_001091992.1) encode the same isoform.

Immunogen

Peptide with sequence AETEALSKLREDFR-C, from the N Terminus of the protein sequence according to NP_001091990.1; NP_006401.3; NP_001091991.1; NP_001091992.1; NP_001091993.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 28kDa band observed in lysates of cell line HeLa (calculated MW of 27.0kDa according to NP_006401.3). Recommended concentration: 0.3-1µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB06323 (0.6 μ g/ml) staining of HeLa lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.