

### **UK Office**

Everest Biotech Ltd Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD UK

Enquiries: info@everestbiotech.com Sales: sales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

# **US Office**

#### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106 Ramona, CA 92065 USA

Inquiries: info@everestbiotech.com Sales: usasales@everestbiotech.com Tech support: support@everestbiotech.com

Tel: 888-320-4628 (toll-free) Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

### EB08298 - Goat Anti-TMPRSS4 (aa245-57) Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: TMPRSS3, transmembrane protease, serine 4, MT-SP2, membrane-type serine protease 2, transmembrane serine protease 3, type II membrane serine protease Official Symbol: TMPRSS4 Accession Number(s): NP\_063947.1; NP\_001077416.1 Human GeneID(s): <u>56649</u> Important Comments: This antibody is expected to recognise both reported isoforms (NP\_063947.1 and NP\_001077416.1).

#### Immunogen

albumin.

Peptide with sequence HCFRKHTDVFNWK, from the internal region of the protein sequence according to NP\_063947.1; NP\_001077416.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

Aliquot and store at -20°C. Minimize freezing and thawing.


## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:1000.

**Western blot:** Approx. 48kDa band observed in lysates of cell lines HEK293 and A549 (calculated MW of 48.2kDa according to NP\_063947.1). Recommended concentration: 0.3-1µg/ml.

#### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human



EB08298 (1µg/ml) staining of HEK293 lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.