



Everest Biotech Ltd  
Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD, United Kingdom

[everestbiotech.com](http://everestbiotech.com)

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel +44 1869 238326

Fax +44 1869 238327

**Research Use Only. Not for diagnostic or therapeutic use.**

Storage: For long-term storage keep aliquots at -20°C. (Store no longer than 12 months at 4°C). Minimize freezing and thawing.

## EB08298 - Goat Anti-TMPRSS4 (aa245-57) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** type II membrane serine protease, transmembrane serine protease 3, membrane-type serine protease 2, MT-SP2, transmembrane protease, serine 4, TMPRSS3

**Official Symbol:** TMPRSS4

**Accession Number(s):** NP\_063947.1; NP\_001077416.1

**Human GeneID(s):** [56649](#)

**Important Comments:** This antibody is expected to recognise both reported isoforms (NP\_063947.1 and NP\_001077416.1).

### Immunogen

Peptide with sequence HCFRKHTDVFNWK, from the internal region of the protein sequence according to NP\_063947.1; NP\_001077416.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:1000.

**Western blot:** Approx. 48kDa band observed in lysates of cell lines HEK293 and A549 (calculated MW of 48.2kDa according to NP\_063947.1). Recommended concentration: 0.3-1µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB08298 (1 $\mu$ g/ml) staining of HEK293 lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.