



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08297 - Goat Anti-TMPRSS4 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** transmembrane protease, serine 4, MT-SP2, TMPRSS3, membrane-type serine protease 2, transmembrane serine protease 3, type II membrane serine protease

**Official Symbol:** TMPRSS4

**Accession Number(s):** NP\_063947.1; NP\_001077416.1

**Human GeneID(s):** [56649](#)

**Non-Human GeneID(s):** 214523 (mouse)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_063947.1 and NP\_001077416.1).

### Immunogen

Peptide with sequence DSTRCNADDAYQGE, from the internal region of the protein sequence according to NP\_063947.1; NP\_001077416.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 48kDa band observed in lysates of cell line HEK293 (calculated MW of 48.2kDa according to NP\_063947.1 and of 47.6kDa according to NP\_001077416.1). Recommended concentration: 0.5-1.5µg/ml. Primary incubation was 1 hour.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse



EB08297 (0.5µg/ml) staining lysates of cell line HEK293 (35µg protein in RIPA buffer). Detected by chemiluminescence.