

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

Enquiries:

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB06739 - Goat Anti-Tyrosine Hydroxylase Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: TH, tyrosine hydroxylase, HGNC:11782, TYH, DYT14, DYT5b,

dystonia 14, tyrosine 3-monooxygenase

Official Symbol: TH

Accession Number(s): NP_954986.2; NP_000351.2; NP_954987.2

Human GeneID(s): 7054

Important Comments: This antibody is expected to recognise all three reported isoforms

(as represented by NP_954986.2; NP_000351.2; NP_954987.2).

Immunogen

Peptide with sequence C-VQDELDTLAHAL, from the C Terminus of the protein sequence according to NP_954986.2; NP_000351.2; NP_954987.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:8000.


IHC: Cryosections of Human Hypothalamus. Recommended dilution range between

1:1000 and 1:10.000.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Rat, Dog



EB06739 immunostaining of dopaminergic neurons in cryosection of an immersion-fixed (4% PFA) human hypothalamus. HRP-staining with Ni-DAB, after Biotin-SP-anti-goat (IgG) method. Data obtained by Drs. Éva Rumpler and Erik Hrabovszky, Inst, Exp, Med, Budapest, Hungary.