

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona,

CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB10535 - Goat Anti-UGT8 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: UGT8, UDP glycosyltransferase 8, CGT, UGT4,

2-hydroxyacylsphingosine 1-beta-galactosyltransferase, UDP-galactose ceramide galactosyltransferase, UDP-galactose-ceramide galactosyltransferase, ceramide

UDP-galactosyltransferase, cerebroside synthase

Official Symbol: UGT8

Accession Number(s): NP_003351.2

Human GeneID(s): 7368

Non-Human GenelD(s): 50555 (rat)

Important Comments: Reported variants represent identical protein: NP_001121646.1,

NP_003351.2.

Immunogen

Peptide with sequence C-NGKYKRNGHIK, from the internal region of the protein sequence according to NP_003351.2.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: In transfected MCF7 transiently expressing Human UGT8 a band of approx. 65kDa is observed. This band is not observed in the non-transfected MCF7. The calculated molecular size is 61.5kDa according to NP_003351.2. Recommended concentration, 1-3µg/ml. Preliminary experiments gave an approx 50kDa band in Human Brain lysates and 48KDa in Rat Brain lysates after 1µg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 61.5kDa according to NP_003351.2. The observed band was successfully blocked by incubation with the immunizing peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Rat, Dog, Cow



MCF7 overexpressing Human UGT8 and probed with EB10535 (mock transfection in first lane). Primary incubation (1ug/ml) was overnight at 4C. Data provided by Tomasz Owczarek, Wrocław University, Poland.

Detected by chemil