

GOAT ANTI-58K GOLGI PROTEIN(N-TERM)/FTCD, BIOTINYLATED ANTIBODY

SKU: EB06442-B



SPECIFICATIONS

| | |
|------------------------------|---|
| Formulation | Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. |
| Unit Size | 100 µg |
| Storage Instructions | Aliquot and store at -20°C. Minimize freezing and thawing. |
| Synonym / Alias Names | formiminotransferase-cyclodeaminase formiminotransferase cyclodeaminase LCHC1 formimidoyltransferase cyclodeaminase FTCD |
| Accession ID | NP_006648.1 |
| Blocking Peptide | EBP06442-B |
| Immunogen | Peptide with sequence SQLVECVPNFSEGKNQ., from the N Terminus of the protein sequence according to NP_006648.1. |
| Product Comments | Reported variants represent identical protein: NP_996848.1, NP_006648.1 Reported variants represent identical protein: XP_006724024.2, XP_011527739.1 |
| Peptide Sequence | SQLVECVPNFSEGKNQ. |
| Purification Method | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Shipping Instructions | Refrigerated |
| Predicted Species | Human, Mouse, Pig |
| Reactive Species | Human |
| Human Gene ID | 10841 |
| Mouse Gene ID | 14317 |
| Product Grade | https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png |
| ELISA Detection Limit | Antibody detection limit dilution 1:64000. |
| Western Blot | Approx 58kDa band observed in Human Liver lysates (calculated MW of 58.9kDa according to NP_006648.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 1-3µg/ml. |
| Application Type | Pep-ELISA, WB |

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

