

GOAT ANTI-AID ANTIBODY

SKU: EB08451

250kDa
150kDa
100kDa
75kDa
50kDa

37kDa

25kDa

20kDa

15kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage	Aliquot and store at -20°C. Minimize freezing and thawing.
Instructions	
Synonym /	integrated into Burkitt\ls lymphoma line Ramos activation-induced deaminase HIGM2 CDA2 ARP2 activation-
Alias	induced cytidine deaminase AICDA AID
Names	
Accession ID	NP_065712.1
Blocking Peptide	EBP08451
Immunogen	Peptide with sequence RRKFLYQFKKNVR-C, from the N Terminus of the protein sequence according to NP_065712.1.
Peptide Sequence	RRKFLYQFKKNVR-C
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Pig, Cow
Reactive Species	Human
Human Gene ID	57379
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA	
Detection Limit	Antibody detection limit dilution 1:1000.
Western Blot	Approx 26kDa band observed in Human Tonsil lysates (calculated MW of 24.0kDa according to NP_065712.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.
Application Type	Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa