

Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-APOBEC3G ANTIBODY

SKU: EB07744







250kDa 150kDa 100kDa

75kDa 50kDa

37kDa

25kDa 20kDa

15kDa





Telephone: (650) 697-3600

SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 μg

Storage

Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

phorbolin-like protein MDS019|apolipoprotein B mRNA editing enzyme catalytic polypeptide-like

Synonym / 3G|OTTHUMP00000028911|DNA dC->dU editing **Alias**

enzyme|dJ494G10.1|bK150C2.7|MDS019|FLJ12740|CEM15|ARP9|apolipoprotein B mRNA editing enzyme, **Names**

catalytic polypeptide-like 3G|APOBEC3G

Accession

NP 068594.1

Blocking

EBP07744 **Peptide**

Peptide with sequence C-DEHSQDLSGRLR, from the C Terminus of the protein sequence according to **Immunogen**

NP 068594.1.

Peptide

C-DEHSQDLSGRLR **Sequence**

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

Method using the immunizing peptide.

Shipping

Refrigerated Instructions

Predicted

Human **Species**

Reactive

Human **Species**

Human

Gene ID

60489

Product Grade

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png

IHC Results Paraffin embedded Human Testis and Kidney. Recommended concentration: 2.5µg/ml.

ELISA

Detection Antibody detection limit dilution 1:16000.

Limit

Western Approx. 45kDa band observed in lysates of cell line Daudi (calculated MW of 46.4kDa according to

NP_068594.1). Recommended concentration: 0.5-1.5µg/ml. Primary incubation was 1 hour. Blot

Application

Pep-ELISA, WB, IHC **Type**

GALLERY IMAGES



Telephone: (650) 697-3600



250kDa 150kDa 100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa



