


Email: customerservice@vectorlabs.com

Telephone: (650) 697-3600

GOAT ANTI-ARHGDIB ANTIBODY





SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 μg

Instructions Aliquot and store at -20°C. Minimize freezing and thawing. Storage

Synonym /

Names

ARHGDIB|RhoGDI2|Rho GDP-dissociation inhibitor 2|Rho GDP dissociation inhibitor (GDI) beta|Rho GDI

Alias 2|RAP1GN1|Ly-GDI|LYGDI|GDID4|GDIA2|D4

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of A431 cells. Usage Recommended concentration: 10µg/ml. Flow Cytometry: Flow cytometric analysis of A431

Summary cells. Recommended concentration: 10ug/ml.

Accession

NP 001166.3

Blocking

EBP10223 **Peptide**

Immunogen Peptide with sequence TEKAPEPHVE-C, from the N Terminus of the protein sequence according to NP_001166.3.









Peptide TEKAPEPHVE-C

Sequence

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using

Method the immunizing peptide.

Shipping Refrigerated Instructions

Predicted

Human, Cow **Species**

Reactive

Human **Species**

Human

397

Gene ID Product

https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite plus medium.png Grade

ELISA

Detection Antibody detection limit dilution 1:4000.

Limit

Western

Approx. 25kDa band observed in lysates of cell lines Daudi, Jurkat, K562, U937, peripheral blood Monocytes (PBM) and in Human Thymus and Cerebellum lysates, and approx. 26kDa in lysates of cell line MOLT4 (calculated MW of 23.0kDa according to NP_001166.3). Recommended concentration: 0.01-3µg/ml. Primary incubation 1 hour at room

temperature.


Application


Type


Blot

Pep-ELISA, WB, IF, FC

GALLERY IMAGES











Telephone: (650) 697-3600



