

GOAT ANTI-BAG2 ANTIBODY

SKU: EB05167

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

10kDa

SPECIFICATIONS

Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage Instructions	Aliquot and store at -20°C. Minimize freezing and thawing.
Synonym /	dJ417I1.2 (BAG-family molecular chaperone regulator 2) dJ417I1.2 MGC149462 KIAA0576 RP3-496N17.2 BAG-
Alias	family molecular chaperone regulator-2 BAG-2 BCL2-associated athanogene 2 BAG2
Names	
Accession ID	NP_004273.1
Blocking Peptide	EBP05167
Immunogen	Peptide with sequence CSKTLQQNAESRFN, from the C Terminus of the protein sequence according to NP_004273.1.
Peptide Sequence	CSKTLQQNAESRFN
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Pig, Cow
Reactive Species	Human, Pig
Human Gene ID	9532
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:8000.
Western Blot	Approx 25kDa band observed in Human HeLa lysates (calculated MW of 23.7kDa according to NP_004273.1). Recommended concentration: 0.1-0.3µg/ml.
Application Type	Pep-ELISA, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa

10kDa

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

15kDa