

GOAT ANTI-BCL7A ANTIBODY

SKU: EB05006

250kDa

150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

SPECIFICATIONS


Formulation	Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.
Unit Size	100 µg
Storage Instructions	Aliquot and store at -20°C. Minimize freezing and thawing.
Synonym /	
Alias	B-cell CLL/lymphoma-7 B-cell CLL/lymphoma 7A BCL7 BCL7A
Names	
Usage Summary	Immunofluorescence: Strong expression of the protein seen in the nuclei of U2OS cells. Recommended concentration: 10µg/ml. <p>Flow Cytometry: Flow cytometric analysis of Daudi cells. Recommended concentration: 10ug/ml.</p>
Accession ID	NP_066273.1; NP_001019979.1
Blocking Peptide	EBP05006
Immunogen	Peptide with sequence C-MKLEASQQNSEEM, from the C Terminus of the protein sequence according to NP_066273.1; NP_001019979.1.
Product Comments	This antibody is expected to recognize both reported isoforms (NP_066273.1; NP_001019979.1).
Peptide Sequence	C-MKLEASQQNSEEM
Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Mouse, Rat, Dog, Cow, Pig
Reactive Species	Human
Human Gene ID	605
Mouse Gene ID	77045
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_plus_medium.png
ELISA Detection Limit	Antibody detection limit dilution 1:4000.
Western Blot	Approx 26kDa band observed in Human Lymph Node lysates (calculated MW of 25.0kDa according to NP_066273.1). Recommended concentration: 1-3µg/ml. Primary incubation 1 hour at room temperature.
Application Type	Pep-ELISA, WB, IF, FC

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES

250kDa



150kDa

100kDa

75kDa

50kDa

37kDa

25kDa

20kDa

